arXiv:1412.5092v1 [math.FA] 15 Dec 2014

# Rigged Hilbert spaces and inductive limits

S.A. Pol'shin*

**Abstract**

We construct a nuclear space $\Phi$ as an inductive limit of finite-dimensional subspaces of a Hilbert space $H$ in such a way that $(\Phi, H, \Phi')$ becomes a rigged Hilbert space, thus simplifying the construction by Bellomonte and Trapani.

MSC 2010: 47A70 46A13

Keywords: rigged Hilbert space, inductive limit, LB-space

Rigged Hilbert spaces (RHS) introduced by Gelfand and Kostyuchenko in 1955 now play an important role in the theory of eigenfunction expansions [11, 3, 13] as well as in the theory of resonant states in quantum mechanics (see [5, 7] for a review). Constructions of rigged Hilbert spaces used in physics are based on spaces of analytic functions [10, 6]. Recently the use of inductive limits was proposed to construct RHS [2]. In the present note we simplify the method of [2] by restricting ourselves to strict inductive limits of spectra of finite-dimensional Hilbert spaces. Also we fill the gap in the construction presented in [12].

Recall [8] that a *direct spectrum* is a family $\{H_i; i \in \mathbb{N}\}$ of LCVS together with continuous maps $\varphi_{ij} : H_i \to H_j$ for all $i < j$ such that the equality $\varphi_{ij}\varphi_{jk} = \varphi_{ik}$ holds for all $i < j < k$.

Suppose $\{H_i\}$ is an inflating sequence of LCVS (i.e. $H_i \subset H_{i+1}$ for all $i$), then $\Phi = \bigcup_{n-1}^{\infty} H_i$ endowed with inductive topology w.r.t. canonical embeddings $\varphi_i : H_i \to \Phi$ is called *inductive limit* of the direct spectrum $\{H_i\}$ denoted by $\lim_{\to} H_i$. Then due to well-known universality of inductive topology ([14], II.6.1; [4], 2.3.5) we have analogous universality property of direct limit [8].

**Proposition 1.** *Let $\tilde{\Phi}$ be a LCVS and let $\tilde{\varphi}_i : H_i \to \tilde{\Phi}$ be continuous maps such that $\varphi_{i\,i+1}\tilde{\varphi}_i = \tilde{\varphi}_{i+1}$ for all $i$. Then the continuous map $\lambda : \Phi \to \tilde{\Phi}$ there exists such that the diagram*




(1)

*commutes, where $\Phi = \lim_{\to} H_i$.*

Taking $\tilde{\Phi} = \mathbb{C}$ in Prop. 1 we obtain the following corollary due to Dieudonné and Schwartz.

*polshin.s@gmail.com

**Corollary 1.** *Any linear functional on* $\lim_{\to} H_i$ *is continuous iff its restrictions to each* $H_i$ *are.*

Suppose now that $H_i \neq H_{i+1}$ for all $i$ and the topology of $H_i$ is equivalent to those induced from $H_{i+1}$, then the inductive limit is called *strict*. It is well known that the strict inductive limit is complete, Hausdorff and nuclear provided all the $H_i$ are ([14], II.6.4, II.6.6, III.7.4; [4], 2.6.2, 2.6.5).

**Definition 1.** Let $i: \Phi \to H$ be an injective homomorphism of complete nuclear space $\Phi$ into Hilbert space $H$ with dense range. Then $(\Phi, H, \Phi')$ is called *Gelfand triple* (or *RHS*).

**Example 1.** Consider the construction of [12] based on the embedding of the space $s$ of rapidly decreasing sequences into $\ell^2$. The range of this embedding is indeed dense in $\ell^2$ (each finite-dimensional subspace of $\ell^2$ contains an element of $s$). It remains to show that this embedding is continuous, but it follows from the following definition of topology of $s$ by seminorms equivalent to ordinary one ([4], 1.3.19):

$$q_k(x)^2 = \sum_{n=1}^{\infty} n^{4k} |x_n|^2.$$

In particular, we can take functional realization $s \cong C^\infty(\mathbf{T})$ and $\ell^2 \cong L^2(\mathbf{T})$.

Let $H \cong \ell^2$ be a separable infinite-dimensional Hilbert space with basis $\{e_i, i \in \mathbb{N}\}$, then for any $x = \sum_{i=1}^{\infty} x_i e_i \in H$ we can define its projections $P_n x = \sum_{i=1}^{n} x_i e_i \in H_n = \operatorname{span}\{e_i, i = 1, \dots, n\}$. Let $\{\alpha_i, i \in \mathbb{N}\}$ be an increasing sequence of natural numbers, we see that $\Phi = \lim_{\to} H_{\alpha_i}$ is dense in $H$ since $\|x - P_n x\|_H \to_{n\to\infty} 0$. Then from Prop. 1 it follows that the continuous map $\lambda: \Phi \to H$ there exist and it is injective. Indeed, otherwise we have $\lambda \varphi_{\alpha_i}(x) = 0$ for some $i$ and $x \neq 0$ in $H_{\alpha_i}$, but this violates the injectivity of $\tilde{\varphi}_i$ in the diagram (1).

The above construction may be reversed in the following sense. Let $\Phi = \lim_{\to} H_i$ be a strict inductive limit of a sequence of *finite-dimensional* Hilbert spaces. Consider the bilinear form on $\Phi$ defined by $\langle x, y\rangle_\Phi = \langle x, y\rangle_{H_i}$ for $x, y \in H_i$. This definition is correct since all the $\varphi_{ij}$ are isometries and $\langle -, -\rangle_\Phi$ is continuous due to Corollary 1. Then we can take Hilbert space $H$ to be (isomorphic to) the completion of $\Phi$ w.r.t. this norm in the spirit of original construction presented in [11]; see also [9] for another definition of continuous norm on inductive limit spaces. So we have proved the following proposition.

**Proposition 2.** *Consider the following data:*

- *Separable infinite-dimensional Hilbert space* $H$.
- *Strict inductive limit of a sequence of finite-dimensional Hilbert spaces.*

*Then starting from one of these data we can construct another one in such a way that* $(\Phi, H, \Phi')$ *becomes a RHS.*

**Example 2.** Let $H_n$ be the space of polynomials on $\xi \in \mathbf{R}$ on degree $\leq n$ and let $H = L^2(\mathbf{R})$. Consider continuous embedding $H_n \to H: P(\xi) \mapsto e^{-x^2/4} P(\xi)$, then due to the density theorem ([1], Theorem 4.5.1) we see that $\Phi = \lim_{\to} H_n$ is dense in $H$, so $(\Phi, H, \Phi')$ is a RHS.

# References


[1] P. Antosik, J. Mikusinski and R. Sikorski, *Theory of distributions. The sequential approach,* Amsterdam, Elsevier; Warszawa, PWN, 1973.

[2] G. Bellomonte and C. Trapani, *Rigged Hilbert spaces and contractive families of Hilbert spaces,* Monatsh. Math. 164 (2011), 271-285.

[3] Yu.M. Berezanskiĭ, *Expansions in eigenfunctions of selfadjoint operators,* Translations of Mathematical Monographs 17, Providence, American Mathematical Society, 1968.

[4] V.I. Bogachev, O.G. Smolyanov and V.I. Smirnov *Topological vector spaces and their applications*, R&C Dynamics, Moscow-Izhevsk, 2012. (in Russian)

[5] A. Böhm and M. Gadella, *Dirac kets, Gamov vectors and Gelfand triplets. The rigged Hilbert space and quantum mechanics*, Lecture Notes in Physics 78, Berlin, Springer-Verlag, 1978.

[6] C.G. Bollini, O. Civitarese, A.L. De Paoli and M.C. Rocca, *Gamow states as continuous linear functionals over analytical test functions*, J. Math. Phys. 37 (1996), 4235-4242.

[7] O. Civitarese and M. Gadella, *Physical and mathematical aspects of Gamow states,* Physics Reports 396 (2004), 41-113.

[8] K. Floret, *Some aspects of the theory of locally convex inductive limits.* in: Functional analysis: surveys and recent results, II, Bierstedt K.-D. and Fuchssteiner B. (eds.), North-Holland Math. Stud. 38, North-Holland, Amsterdam-New York, 1980, 205-237.

[9] K. Floret, *Continuous norms on locally convex strict inductive limit spaces,* Math. Z. 188 (1984), 75-88.

[10] M. Gadella, *A rigged Hilbert space of Hardy-class functions: Applications to resonances.* J. Math. Phys. 24 (1983), 1462-1469.

[11] I.M. Gel'fand and N.Ya. Vilenkin, *Generalized functions.*, Vol. 4, Academic Press, New York - London, 1964.

[12] G. Lindblad and B. Nagel, *Continuous bases for unitary irreducible representations of* $SU(1,1)$, Ann. Inst. H. Poincaré, Sect. A (N.S.) 13 (1970), 27-56.

[13] K. Maurin, *General eigenfunction expansions and unitary representations of topological groups,* Monografie Matematyczne 48, Warszawa, PWN, 1968.

[14] H. Schaefer, *Topological vector spaces*, Macmillan, New York, 1966.



Institute for Theoretical Physics,
NSC Kharkov Institute of Physics and Technology,
Akademicheskaia St. 1, 61108 Kharkov, Ukraine